

## ANTIOXIDANT AND ANTIPROLIFERATIVE ACTIVITY OF SMALL PEPTIDES ISOLATED FROM MARINE ALGAE BY GREEN METHODS

Elena MIHAI, Alexandra GASPAR-PINTILIESCU, Laura Mihaela STEFAN,  
Rodica TATIA, Viorica COROIU, Oana CRACIUNESCU

National Institute of Research and Development for Biological Sciences,  
296 Splaiul Independenței, Bucharest, Romania

Corresponding author email: elena.mihai@incdsb.ro

### Abstract

*This study aimed to investigate new green methods based on combined ultrasound- and enzyme-assisted technologies to isolate bioactive peptides from Cladophora vagabunda green macroalga. Also, their antioxidant and antiproliferative activity was analysed in view of biomedical applications. Algal powder was sonicated in alkaline buffer, pH 8, at 37 °C, for 1 h. The extract was treated with proteinase K and alcalase, respectively, to obtain two protein hydrolysates. Each hydrolysate was fractionated by centrifugal ultrafiltration using filter membranes with molecular weight cut-off (MWCO) at 3 kDa, to separate small peptides. The fractions were analyzed for Trolox Equivalent Antioxidant Capacity (TEAC). The results showed higher capacity of hydrolysates obtained by alcalase treatment to scavenge free radicals. Additionally, the antiproliferative activity was evaluated in HT-29 tumor cells cultivated in the presence of peptides obtained by alcalase treatment and the results showed a decrease of cell viability below 70% after 48 h of cultivation. In conclusion, peptide fractions isolated by green techniques from C. vagabunda alga had significant biological activity and are recommended for further testing as therapeutic agents.*

**Key words:** algal peptides, Cladophora vagabunda, enzymatic hydrolysis, marine bioactive compounds, sonication.

### INTRODUCTION

High quantities of animal proteins are found in sources such as meat, eggs and milk, while vegetable proteins are found in soy and maize. Lately, the micro- and macroalgal biomass was considered a viable alternative source of proteins (Espinosa-Ramirez et al., 2023). The protein production from marine macrophyte algae (2.5-7.5 tons/ha/year) is higher than that from terrestrial cultures of soy (0.6-1.2 tons/ha/year) (O'Connor et al., 2022). Additionally, proteins from edible seaweeds are nutritionally important, containing high quantities of essential amino acids like lysine, tryptophan, methionine, valine, thus being recommended for human diet (Kazir et al., 2019). The protein extraction yield represented up to 47% when using red algae species (Rhodophyceae) and up to 29% in green (Chlorophyceae) and brown (Phaeophyceae) algae species, the quantity varying according to geographical area and harvesting season (Vieira et al., 2018). However, traditional technologies for algal protein extraction are time consuming

and economically unviable (Bleakley & Hayes, 2017).

Currently developed technologies for bioactive compounds extraction from algal biomass are based on green methods of ultrasound-assisted extraction (UAE), microwave-assisted extraction (MAE), and enzyme-assisted extraction (EAE). In case of UAE, the effect of acoustic cavitation could improve seaweed proteins bioavailability by formation and growth of vapor bubbles and facilitating penetration of solvent into the cells to release intracellular compounds (Arya et al., 2023). An ultrasound-assisted treatment in acid or alkaline conditions was efficient for protein extraction from *Ascophyllum nodosum* brown alga (Pan-utai et al., 2023). On the other side, MAE takes place by transfer of microwaves energy to polar water molecules, increasing their vibration, collision, and intracellular heating, and thus leading to an effect of pressurization and algal cell membrane damage (Grosso et al., 2015). EAE using enzymes (cellulases, agarases, xylanase, carragenase) for digestion of cellulose-type polysaccharides (galactans, xylans, fucoidans, laminarin, alginates, carrageenan) present in the

rigid cell wall of algal cells were useful in protein extraction from red alga species *Chondrus crispus*, *Gracilaria verrucosa* and *Palmaria palmata* (Fleurence et al., 1995).

Green algae present a chemical composition rich in polysaccharides, phenolic compounds, proteins and amino acids, fatty acids, vitamins and minerals (Lafarga et al., 2020). Extracts of various green algae demonstrated immunomodulatory, antimicrobial, anti-inflammatory and antitumoral activity (Surayot et al., 2016). The genus *Cladophora* is comprising marine green macrophyte algae with filamentous branched multinucleate cells, widespread from temperate to arctic and tropical waters (Michalak & Messyas, 2020). *Cladophora vagabunda* is abundantly found in the Black Sea, it has an invasive tendency, but it is still an underexploited biological resource (Marin & Timofte, 2011). *Cladophora* biomass was reported as valuable raw material for agriculture and cosmetics (Messyas et al., 2015) and extraction of volatile compounds and fatty acids from *C. vagabunda* indicated applications as bioadditives and biopreservatives (Horincar et al., 2014). In this context, the present paper aimed to investigate new green methods for the isolation of bioactive peptides from *C. vagabunda* macroalga and their antioxidant and antiproliferative activity, in view of valorization as natural ingredients with biomedical applications.

## MATERIALS AND METHODS

### Biological material and chemical reagents

The biological material consisted of *C. vagabunda* green macrophyte alga collected from the Black Sea between 2 Mai and Vama Veche, in August 2022. The algae were extensively washed in cold tap water to remove the sand, dried in an oven (Memmert, Germany) at 30 °C and ground using an electric grinder until a fine green powder was obtained.

All chemical reagents used in this study were of analytical grade. 2,4,6-trinitrobenzene sulfonic acid (TNBS), 2,2'-azino-bis (3-ethyl-benzothiazoline-6-sulfonic acid (ABTS), proteinase K (E.C. 3.4.21.64), alcalase (E.C. 3.4.21.14), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) and all other chemicals were purchased from Sigma-Aldrich (Germany), unless otherwise stated. Human HT-29

epithelial cell line derived from colorectal adenocarcinoma, Dulbecco's Modified Eagle Medium (DMEM), fetal calf serum (FCS) and penicillin-streptomycin-neomycin antibiotic mixture (PSN) were purchased from Sigma-Aldrich (Germany).

### Extraction of algal proteins based on ultrasound-assisted technology

The extraction of proteins from *C. vagabunda* powder was carried out in TES buffer 0.05 M, pH 8 (1:20, m/v) by ultrasonication in a sonication bath (Elma, Germany), at 60 kHz, at a temperature of 37 °C, for 1 h (Figure 1). The temperature was controlled throughout the process not to exceed 37 °C, by sequencing the ultrasonication process in 5 on-off cycles, each lasting 10 min with a break of 2 min. Then, the process continued by magnetic stirring, at 4 °C, for 24 h. The extraction was repeated twice using the residue obtained after centrifugation of the solution at 9000 g, at 4 °C, for 20 min, and all supernatants were pooled together (P1).

The extraction yield was determined on a dry weight (d.w.) basis using the following formula:

$$\text{extraction yield (\%)} = \text{d.w.}_{\text{final}} / \text{d.w.}_{\text{initial}} \times 100$$

The protein content was determined by BCA assay, as previously described (Mihai et al., 2021).

### Preparation and purification of algal peptides by enzyme-assisted technology

Enzymatic hydrolysis of protein extract (P1) was carried out using specific neutral proteases of microbial origin under specific conditions of pH and temperature. Thus, the algal extract aliquots were treated with 3% proteinase K in 0.05 M Tris buffer, pH=8, supplemented with 1 mM CaCl<sub>2</sub> (PP1) and 3% alcalase in 0.05 M Tris buffer, pH=8 (PA1), respectively (Figure 1). The mixtures were incubated in a shaking water bath (Witeg, Germany), at 55 °C, for 3 h. The pH of the reaction mixture was periodically verified throughout the enzymatic process and maintained at the initial optimal value. At the end of the incubation period, the mixtures were heated at 95 °C for 5 min, to inactivate the enzyme. Then, the solutions were cooled at room temperature and centrifuged at 9000 g, at 4 °C, for 20 min. Each hydrolysate was

subjected to centrifugal ultrafiltration using cellulose membrane filter units with MWCO at 3 kDa (Amicon, Germany) centrifuged at 7500 g in successive cycles of 30 min, according to manufacturer's instructions. Thus, two fractions were separated from each hydrolysate, the permeates (PP1 and PA1 fractions with MW<3 kDa) and the retentates (PP1 and PA1 fractions with MW>3 kDa).

#### **Determination of the degree of hydrolysis**

The degree of hydrolysis was determined as free amino groups quantified by TNBS assay, as previously described (Craciunescu et al., 2011). Thus, samples of hydrolysates (PP1, PA1) and the non-hydrolysed protein extract (P1) (0.25 ml) were mixed with 0.05% TNBS in 0.05 M TES buffer, pH=8 (1 ml) and incubated in a shaking water bath (Witeg, Germany) at 50 °C, in the dark, for 1 h. Then, 0.1 N HCl (3 mL) was added and the incubation continued in the water bath, at 50 °C, in the dark, for 10 min. After cooling at room temperature, distilled water (5 mL) was added. The absorbance (Abs) was read in quartz cuvettes at a wavelength of 346 nm using a V-650 UV-VIS spectrophotometer (Jasco, Japan). Total hydrolyzate was prepared by P1 incubation in 6 N HCl, at 110 °C, 24 h. The degree of hydrolysis was calculated using the following formula:

$$\text{degree of hydrolysis (\%)} = (\text{Abs}_{\text{hydrolysate}} - \text{Abs}_{\text{non-hydrolysed}}) / (\text{Abs}_{\text{total hydrolysate}} - \text{Abs}_{\text{non-hydrolysed}}) \times 100$$

#### **Determination of antioxidant capacity**

The antioxidant capacity was determined by TEAC assay, as previously described (Gaspar-Pintilieșcu et al., 2022). A stock solution was prepared by mixing 7 mM ABTS with 2.45 mM potassium persulfate (1:1, v/v) and incubation at room temperature, in the dark, for 16 h, to allow free ABTS radicals formation. For the experiment, the stock solution was diluted with distilled water to give an Abs of  $0.70 \pm 0.02$  at a wavelength of 734 nm (blank). The samples (100 μL) were mixed with ABTS reagent (1 mL) and incubated at room temperature, in the dark, for 10 min. The Abs of the reaction mixtures was read at 734 nm using a V650 UV-VIS spectrophotometer (Jasco, Japan). A standard curve was built using Trolox, a synthetic

antioxidant, in the range of concentrations of 10-150 μM. The antioxidant capacity was calculated using the following formula:

$$\text{TEAC} = (\text{Abs}_{\text{blank}} - \text{Abs}_{\text{sample}}) / \text{Abs}_{\text{blank}} \times 100$$

The results were expressed as mM Trolox equivalents (TE) per g d.w.

#### **Determination of antiproliferative activity**

##### **Cell culture and treatment**

Human tumor HT-29 epithelial cells were seeded in the wells of a 96-well culture plate at a density of  $5 \times 10^4$  cells/mL and cultured in DMEM medium supplemented with 10% FCS, 1% glutamine, and 1% PSN antibiotic mixture. The plates were incubated in standard conditions, at a temperature of 37 °C, in 5% CO<sub>2</sub> atmosphere, for 24 h, to allow cell adhesion. Then, the culture medium was replaced with fresh culture medium supplemented with FCS containing different concentrations of algal peptides in the concentration range of 0.16-2.5 mg/mL. The plates were incubated in standard conditions for 24 and 48 h, respectively. The cells cultivated in MEM medium without treatment, in standard conditions served as negative control.

##### **MTT assay**

Cell proliferation was evaluated by MTT assay based on the reduction of yellow tetrazolium salts to purple insoluble formazan by succinate mitochondrial dehydrogenases in metabolically viable cells, a reaction mediated by the reduction NADH to NADPH (Oprita et al., 2008). Thus, at the end of the incubation period, the culture medium in each well was replaced with 0.25 mg/mL MTT solution (100 μL) and the plates were incubated at 37 °C, in 5% CO<sub>2</sub> atmosphere, for 3 h. Then, the MTT solution was replaced with isopropanol (100 μL) to solubilize the formazan crystals by gentle stirring, on a horizontal shaker, for 15 min. The Abs was read at a wavelength of 570 nm using a SpectroStar Nano microplate reader (BMG Labtech, Germany). The Abs values were directly proportional to the number of viable cells. Cell proliferation was calculated using the following formula:

$$\text{cell proliferation (\%)} = \text{Abs}_{\text{sample}} / \text{Abs}_{\text{control}} \times 100$$

## Statistical analysis

The experiments were performed in triplicate and the results were expressed as mean  $\pm$  standard deviation (SD) (n=3). Statistical analysis was performed by two-tailed, two-sample equal variance Student *t*-test on control-sample pairs of interest. Differences were considered statistically significant at  $p < 0.05$ .

## RESULTS AND DISCUSSIONS

In the present study, an experimental protocol based on combined green technologies of UAE and EAE was optimized at laboratory level for sustainable valorization of *C. vagabunda* green macrophyte algal biomass to obtain peptide fractions with bioactivity towards free radicals scavenging and antiproliferative activity in tumor cell culture, in order to be used as natural therapeutic agents (Figure 1).




Figure 1. Scheme of combined UAE and EAE technologies to isolate algal bioactive peptides under controlled conditions from *C. vagabunda*

### Preparation of algal proteins by UAE

First, an algal protein extract was obtained from *C. vagabunda* green alga powder by ultrasonication with a yield of 23.68%,

calculated on a d.w. basis. This value was similar to data reported in previous studies on different green algae from *Ulva* species (7-29%) (Pimentel et al., 2019).

This high extraction rate confirmed the efficacy of the sonication-based optimized technology. It is known that macrophyte algae are characterized by the presence of polysaccharidic cell walls that play an important role in mechanical shear resistance (Shao & Duan, 2022). As a result, in the present study, the extraction of proteins from *C. vagabunda* green macroalga was conducted by ultrasonication treatment at low frequency and high intensity (20-100 kHz), which allowed the destruction of cell walls for improving the bioactive compounds extraction yield. Additionally, the ultrasound-assisted technology applied for a short processing period allowed extraction of thermolabile proteins without damaging their structure and properties, in accordance to previous reports (Ojha et al., 2020; Prandi et al., 2022). The alkaline buffer (pH=8) improved the solubilization of large peptides and proteins from cell cytoplasm and break down of their globular structure, but also facilitated the easy release of small peptides and free amino acids from algal biomass. Similar studies reported that the yield of protein extraction from *Nannochloropsis* spp. was doubled in alkaline solvents (pH=8), compared to water (O'Connor et al., 2022; Parniakov et al., 2015).

### Preparation of algal peptides by EAE

Algal peptides were obtained using specific neutral proteases (3% proteinase K and 3% alcalase solutions) with broad action and stability over large pH and temperature domains, to ensure controlled degradation (Figure 1). An indicator of protein degradation and peptides formation is the number of free amino groups, determined in the present study by TNBS assay. The degree of enzymatic hydrolysis of algal proteins reached values of 9.22% under proteinase K treatment (PP1) and 45.87% under alcalase treatment (PA1). These data indicated an efficient hydrolysis of *C. vagabunda* protein extract by alcalase under mild optimal reaction conditions (pH 8, 55 °C, 3 h).

Each hydrolysate was subjected to centrifugal ultrafiltration through membranes with MWCO of 3 kDa, which allowed separation of 2 fractions (permeates with MW < 3 kDa and

retentates with MW>3 kDa). The results of protein content analysis in permeates and retentates are given in Table 1.

**Table 1.** Protein content of protein hydrolysate and peptide fractions from *C. vagabunda*

Protein content (mg/g d.w.)	
Proteinase K hydrolysate	
PP1 (total hydrolysate)	54.52 ± 2.74
PP1 < 3 kDa (permeate)	20.10 ± 1.22
PP1 > 3 kDa (retentate)	106.17 ± 7.91
Alcalase hydrolysate	
PA1 (total hydrolysate)	73.91 ± 3.35
PA1 < 3kDa (permeate)	44.52 ± 2.42
PA1 > 3kDa (retentate)	182.49 ± 7.63

The results represent mean ± SD (n=3).

The results showed that alcalase hydrolysate (PA1) had higher protein content (73.91 mg/g d.w.), compared to proteinase K hydrolysate (PP1) (54.52 mg/g d.w.). Also, peptide fractions isolated from alcalase hydrolysate had ~2-fold higher protein content than those isolated from proteinase K hydrolysate (Table 1). All these data allowed selection of alcalase as optimal protease for preparation of algal peptides from *C. vagabunda* and obtained fractions were further investigated for bioactivity *in vitro*.

### Antioxidant capacity of *C. vagabunda* peptide fractions

The high protein content of macrophyte algae and their antioxidant activity is an important reason for separation and valorization in biomedical applications (Surayot et al., 2016). The antioxidant capacity of peptide fractions isolated from *C. vagabunda* green alga by UAE and alcalase treatment was determined by TEAC assay. The results are presented in Table 2.

**Table 2.** Antioxidant capacity of alcalase hydrolysate and peptide fractions from *C. vagabunda*

Sample	TEAC (mM TE/g d.w.)
PA1 (total hydrolysate)	163.78 ± 7.34
PA1<3 kDa (permeate)	123.67 ± 5.91
PA1>3 kDa (retentate)	139.43 ± 4.25

The results represent mean ± SD (n=3).

The results showed that all peptide samples presented high potential to scavenge free ABTS

radicals. It was observed a decrease of the antioxidant capacity of peptides from permeate fraction, compared to alcalase hydrolysate and retentate. This could be due to smaller size of peptides from permeate containing more amino and carboxyl terminal groups, which decreased the antioxidant activity, as previously reported (Zhao et al., 2020).

The antioxidant activity of peptides could be correlated to their MW, but also to the hydrophobic nature of the amino acids present in their sequences. Thus, in the case of VECYGPRPQF peptide isolated from *Chlorella vulgaris* green macrophyte alga, rich in hydrophobic and aromatic amino acids, it was reported a potent antioxidant activity (Sheih et al., 2010). In the present study, the use of alcalase for algal protein hydrolysis could lead to hydrophobic peptides with increased antioxidant activity due to its known activity of preferential cleavage of the protein before aromatic amino acids, compared to proteinase K hydrolysis that had broader specificity and also cleaved before aliphatic residues and glycine residues.

Previous studies on *Palmaria palmata* macrophyte alga identified mycosporine-like amino acids, such as palythine, porphyra-334, shinorine-330, asterin, palythanol, palythene, usujirene, produced as low MW water-soluble secondary metabolites (Yuan et al., 2009). Their structural characteristics consisting of a cyclohexanone or hexenimine core conjugated to the nitrogen of an amino acid allowed significant antioxidant activity (Bedoux et al., 2014). Similar molecules were isolated from *Porphyra umbilicalis* macrophyte alga and used to obtain Helioguard 365® sunscreen, exhibiting antioxidant activity, capacity to absorb UV radiation and dissipate energy in the form of heat (Pandey et al., 2017), limiting the negative effect of fotons on cellular components (Pimentel et al., 2019).

### Antiproliferative activity of *C. vagabunda* peptide fractions

The antiproliferative activity of peptide fractions isolated from *C. vagabunda* green alga was assessed in human tumor HT-29 epithelial cells by MTT assay. The results are presented in Figure 2.

The results showed that alcalase-treated peptide fraction PA1<3 kDa presented similar Abs values at 24 h of cultivation and significantly ( $p<0.05$ ) higher Abs values at 48 h of cultivation, compared to the control, indicating stimulation of cell proliferation. The algal hydrolysate PA1 and peptide fraction PA1>3 kDa could significantly ( $p<0.05$ ) inhibit tumor cell proliferation after 48 h of cultivation, at all tested concentrations, compared to the control.

The highest activity was 2-fold inhibition of cell proliferation at 2.5 mg/mL PA1>3 kDa, after 48 h of cultivation. *In vitro* data demonstrated the antiproliferative activity of algal bioactive peptides isolated from *C. vagabunda* in the experimental model based on human HT-29 colon epithelial cells. These *in vitro* results correlated direct proportional to the antioxidant activity of algal peptides and their MW.




Figure 2. Antiproliferative activity of alcalase hydrolysate (PA1) and peptide fractions from *C. vagabunda* green alga. The results represent mean  $\pm$  SD (n=3). \* $p<0.05$ , compared to control.

Previous studies have reported that two peptides VPGTPKNLDSPR and MPAPSCALPRSVVPPR identified in trypsin hydrolysate of *Porphyra haitanensis* protein extract revealed inhibition of human MCF-7 breast cancer cells and human HepG-2 liver cancer cells proliferation and cell cycle arrest inducing cell apoptosis (Fan et al., 2017). The study on papain hydrolysate of *Pyropia haitanensis* protein extract showed antiproliferation effect of QTDDNHSNVLWAGFSR peptide on human HepG-2 liver cancer cells at a concentration of 500  $\mu$ g/mL that induced low inhibition of human normal liver cells (Mao et al., 2017).

## CONCLUSIONS

Green biotechnologies based on combined UAE and EAE using specific neutral proteases (proteinase K, alcalase) were established to

extract bioactive peptides from *C. vagabunda* green macroalga. Alcalase was selected as optimal enzyme for the isolation of peptide fractions from *C. vagabunda* green macroalga due to higher degree of hydrolysis, compared to proteinase K. The hydrolysate obtained by algal protein digestion using alcalase and peptide fractions isolated after centrifugal ultrafiltration showed significant antioxidant capacity as free ABTS radicals scavenging potential. The experimental model *in vitro* demonstrated that the isolated bioactive peptides could inhibit cell proliferation in human tumor HT-29 epithelial cells.

In conclusion, the algal peptides isolated from *C. vagabunda* had significant biological activity and are recommended for further testing as therapeutic agents.

## ACKNOWLEDGMENTS

We thank prof. dr. Ticuta Negreanu-Pirjol from 'Ovidius' University of Constanta for kind cooperation and providing of the biological material. This research was supported by the Ministry of Research, Innovation and Digitization through Program 1 - Development of the National R&D System, Subprogram 1.2 - Institutional Performance - Projects for Excellence Financing in RDI, contract no. 2PFE/2021, and Program Nucleu, project no. 23020201.

## REFERENCES

- Arya, S.S., More, P.R., Ladole, M.R., Pegu, K., & Pandit, A.B. (2023). Non-thermal, energy efficient hydrodynamic cavitation for food processing, process intensification and extraction of natural bioactives: a review. *Ultrasonics Sonochemistry*, 98, 106504.
- Bedoux, G., Hardouin, K., Marty, C., Taupin, L., Vandanjon, L., & Bourgougnon, N. (2014). Chemical characterization and photoprotective activity measurement of extracts from the red macroalga *Solieria chordalis*. *Botanica Marina*, 57, 291–301.
- Bleakley, S., & Hayes, M. (2017). Algal proteins: extraction, application, and challenges concerning production. *Foods*, 6(5), 33.
- Craciunescu, O., Tardei, Ch., Moldovan, L., & Zarnescu, O., (2011). Magnesium substitution effect on porous scaffolds for bone repair. *Central European Journal of Biology*, 6(3), 301-311.
- Espinosa-Ramirez, J., Mondragon-Portocarrero, A.C., Rodriguez, J.A., Lorenzo, J.M., & Santos, E.M. (2023). Algae as a potential source of protein meat alternatives. *Frontiers in Nutrition*, 10, 1254300.
- Fan, X., Bai, L., Mao, X., & Zhang, X. (2017). Novel peptides with anti-proliferation activity from the *Porphyra haitanensis* hydrolysate. *Process Biochemistry*, 60, 98-107.
- Fleurence, J., Massiani L., Guyader, O., & Mabeau, S. (1995). Use of enzymatic cell wall degradation for improvement of protein extraction from *Chondrus crispus*, *Gracilaria verrucosa* and *Palmaria palmata*. *Journal of Applied Phycology*, 7(4), 393–397.
- Gaspar-Pintilieșcu, A., Mihai, E., Ciucan, T., Popescu, A.F., Luntraru, C., Tomescu, J., & Craciunescu, O. (2022). Antioxidant and acetylcholinesterase inhibition capacity of hydrosols from Lamiaceae plants for biopesticide use: role of phenolics. *International Journal of Food Properties*, 25, 996-1008.
- Grosso, C., Valentão, P., Ferreres, F., & Andrade, P. (2015). Alternative and efficient extraction methods for marine-derived compounds. *Marine Drugs*, 13(5), 3182–3230.
- Horincar, V.B., Parfene, G., Tyagi, A.K., Gottardi, D., Dinica, R., Guerzoni, M.E., & Bahrim, G. (2014). Extraction and characterization of volatile compounds and fatty acids from red and green macroalgae from the Romanian Black Sea in order to obtain valuable bioadditives and biopreservatives. *Journal of Applied Phycology*, 26, 551-559.
- Kazir, M., Abuhassira, Y., Robin, A., Nahor, O., Luo, J., Israel, A., & Livney, Y.D. (2019). Extraction of proteins from two marine macroalgae, *Ulva* sp. and *Gracilaria* sp., for food application, and evaluating digestibility, amino acid composition and antioxidant properties of the protein concentrates. *Food Hydrocolloids*, 87, 194–203.
- Lafarga, T., Acien-Fernandez, F.G., & Garcia-Vaquero, M. (2020). Bioactive peptides and carbohydrates from seaweed for food applications: natural occurrence, isolation, purification, and identification. *Algal Research*, 48, 101909.
- Marin, O., & Timofte, F. (2011). *Atlasul macrofitelor de la litoralul romanesc*, Boldas Publishing House, ISBN 978-606-8066-33-2.
- Messyasz, B., Leska, B., Fabrowska, J., Pikosz, M., Roj, E., Cieslak, A., & Schroeder, G. (2015). Biomass of freshwater *Cladophora* as a raw material for agriculture and the cosmetic industry. *Open Chemistry*, 13, 1108-1118.
- Michalak, I., & Messyasz, B. (2020). Concise review of *Cladophora* spp.: macroalgae of commercial interest. *Journal of Applied Phycology*, 33, 133-166.
- Mihai, E., Oancea, A., Craciunescu, O., Ilie, D., Sanda, C., Balan, D., & Jurcoane, S. (2021). Extraction and identification of the main allergens from fried peanuts. *Scientific Bulletin Series F. Biotechnologies*, XXV(2), 41-46.
- Mao, X., Bai, L., Fan, X., & Zhang, X. (2017). Anti-proliferation peptides from protein hydrolysates of *Pyropia haitanensis*. *Journal of Applied Phycology*, 29, 1623-1633.
- O'Connor, J., Garcia-Vaquero, M., Meaney, S., & Tiwari, B.K. (2022) Bioactive peptides from algae: traditional and novel generation strategies, structure-function relationships, and bioinformatics as predictive tools for bioactivity. *Marine Drugs*, 20, 317.
- Ojha, S.K., Aznar, R., O'Donnell, C., & Tiwari, B.K. (2020). Ultrasound technology for the extraction of biologically active molecules from plant, animal and marine sources. *TrAC Trends in Analytical Chemistry*, 115663.
- Oprita, E.I., Moldovan, L., Craciunescu, O., & Zarnescu, O. (2008). *In vitro* behaviour of osteoblast cells seeded into a COL/β-TCP composite scaffold. *Central European Journal of Biology*, 3(1), 31-37.
- Pandey, A., Pandey, S., Pathak, J., Ahmed, H., Singh, V., & Singh S.P. (2017). Mycosporine-like amino acids (MAAs) profile of two marine red macroalgae, *Gelidium* sp. and *Ceramium* sp. *International Journal of Applied Sciences and Biotechnology*, 5, 12–21.
- Pan-utai, W., Pantoa, T., Roytrakul, S., Praiboon, J., Kosawatpat, P., Tamtin, M., & Thongdang, B. (2023) Ultrasonic-assisted extraction and antioxidant potential of valuable protein from *Ulva rigida* macroalgae. *Life*, 13(1),86.

- Parniakov, O., Barba, F.J., Grimi, N., Marchal, L., Jubeau, S., Lebovka, N., & Vorobiev, E. (2015). Pulsed electric field and pH assisted selective extraction of intracellular components from microalgae *Nannochloropsis*. *Algal Research*, 8, 128–134.
- Pimentel, F.B., Alves, R.C., Harnedy, P.A., FitzGerald, R.J., & Oliveira, M.B.P.P. (2019). Macroalgal-derived protein hydrolysates and bioactive peptides: enzymatic release and potential health enhancing properties. *Trends in Food Science & Technology*, 93, 106-124.
- Prandi, B., Di Massimo, M., Tedeschi, T. et al. (2022). Ultrasound and microwave-assisted extraction of proteins from coffee green beans: effects of process variables on the protein integrity. *Food and Bioprocess Technology*, 15, 2712–2722.
- Shao, Z., & Duan, D. (2022). The cell wall polysaccharides biosynthesis in seaweeds: a molecular perspective. *Frontiers in Plant Science*, 13, 902823.
- Sheih, I.-C., Fang, T.J., Wu, T.-K., & Lin, P.-H. (2010). Anticancer and antioxidant activities of the peptide fraction from algae protein waste. *Journal of Agricultural and Food Chemistry*, 58, 1202–1207.
- Surayot, U., Hun Lee, J., Kanongnuch, C., Peerapornpisal, Y., Park, W., & You, S. (2016). Structural characterization of sulfatedarabinans extracted from *Cladophora glomerata* Kützing and their macrophage activation. *Bioscience, Biotechnology, and Biochemistry*, 80(5), 972-982.
- Vieira, E.F., Soares, C., Machado, S., Correia, M., Ramalhosa, M.J., Oliva-Teles, M.T., & Delerue-Matos C. (2018). Seaweeds from the Portuguese coast as a source of proteinaceous material: Total and free amino acid composition profile. *Food Chemistry*, 269, 264–275.
- Yuan, Y.V., Westcott, N.D., Hu, C., & Kitts, D.D. (2009). Mycosporine-like amino acid composition of the edible red alga, *Palmaria palmata* (dulse) harvested from the west and east coasts of Grand Manan Island, New Brunswick. *Food Chemistry*, 112, 321–328.
- Zhao, Y., Zhao, Q., & Lu, Q. (2020). Purification, structural analysis, and stability of antioxidant peptides from purple wheat bran. *BMC Chemistry*, 14, 58.